
PRODUCT INFORMATION

Syngenta Crop Protection, Inc.

**410 Swing Road
Greensboro, NC 27409**

SUPPLEMENTAL LABEL FOR BRAVO WEATHER STIK®

Fungicide

EPA Reg. No. 50534-188-100

Broad spectrum fungicide for control of plant diseases

Active Ingredient:	
Chlorothalonil (tetrachloroisophthalonitile)	54.0%
Inert Ingredients:	46.0%
Total:	100.0%

Contains 6.0 Pounds of Chlorothalonil per gallon (720 grams per liter)

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This labeling must be in the possession of the user at the time of pesticide application.

Reformulation is prohibited. See individual container labels for repackaging limitations.

DIRECTIONS FOR APPLICATION

Crop	Target Diseases	Use Rate lbs. product/A (lbs. a.i./A)	Remarks
Soybeans Determinate (Southern) Varieties AND Indeterminate (Northern) Varieties	Rust (<i>Phakopsora</i> spp.)	1 3/8 to 2 1/4 pts. (1.0 to 1.7)	Make the first application at the first sign of disease. Alternate with another fungicide registered for soybean rust control. Apply in sufficient water to obtain complete coverage, generally 10-20 gallons water per acre. The minimum re-treatment interval is 14 days. DO NOT exceed total of 3 applications per season. DO NOT apply more than 6 pints Bravo Weather Stik (4.5 lbs. a.i.) per acre during each growing season. DO NOT apply within 6 weeks of harvest. DO NOT feed treated parts to livestock or allow grazing in treated fields.
Beans (Snap)	Rust (<i>Phakopsora</i> spp.)	1 3/8 to 3 pts. (1.0 to 2.25)	Make the first application at the first sign of disease. Alternate with another fungicide registered for bean rust control. Apply in sufficient water to obtain complete coverage, generally 10-20 gallons water per acre. The minimum re-treatment interval is 7 days. DO NOT apply more than 12 pints Bravo Weather Stik (9 lbs. a.i.) per acre during each growing season. DO NOT apply within 7 days of harvest.
Beans (Dry)	Rust (<i>Phakopsora</i> spp.)	1 3/8 to 2 pts. (1.0 to 1.5)	Make the first application at the first sign of disease. Alternate with another fungicide registered for bean rust control. Apply in sufficient water to obtain complete coverage, generally 10-20 gallons water per acre. The minimum re-treatment interval is 7 days. DO NOT apply more than 8 pints Bravo Weather Stik (6 lbs. a.i.) per acre during each growing season. DO NOT apply within 14 days before harvest. Bravo Weather Stik may be applied through sprinkler irrigation equipment. See calibration directions which appear on the product label.

Bravo Weather Stik® trademark of a Syngenta Group Company

©2002 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 50534-188-S4 0802

B-BRA WEA 50534-188-S4- 0802-SL - nsp 10/15/02